



MÀSTER UNIVERSITARI EN OPTOMETRIA I CIÈNCIES DE LA VISIÓ

TREBALL FINAL DE MÀSTER

VISUAL AND MORPHOLOGICAL CHARACTERISTICS IN A PRE-TREATED EXUDATIVE AGE-RELATED MACULAR DEGENERATION POPULATION

ANTÍA SUÁREZ VIDAL

DIRECTORA: Sara Lluch Margarit
CODIRECTORA: María Dolores Merindano Encina
Departament d'Òptica i Optometria

Terrassa, 22 de juny de 2017



MÀSTER UNIVERSITARI EN OPTOMETRIA I CIÈNCIES DE LA VISIÓ

Les Sres. Sara Lluch Margarit i María Dolores Merindano Encina, com a directores del treball

CERTIFIQUEN

Que la Sra. Antía Suárez Vidal ha realitzat sota la seva supervisió el treball **VISUAL AND MORPHOLOGICAL CHARACTERISTICS IN A PRE-TREATED EXUDATIVE AGE-RELATED MACULAR DEGENERATION POPULATION** que es recull en aquesta memòria per optar al títol de màster en optometria i ciències de la visió.

I per a què consti, signem aquest certificat.

Sra Sara Lluch Margarit
Directora del treball

Sra María Dolores Merindano Encina
Directora del treball

Terrassa, 22 de juny de 2017



MÁSTER UNIVERSITARIO EN OPTOMETRIA I CIENCIAS DE LA VISIÓN

CARACTERÍSTICAS VISUALES Y MORFOLÓGICAS EN UNA POBLACIÓN CON DEGENERACIÓN MACULAR ASOCIADA A LA EDAD EXUDATIVA PRE-TRATADA

RESUMEN

La degeneración macular asociada a la edad (DMAE) es la principal causa de ceguera legal en países desarrollados. La DMAE exudativa, en la cual se centra este estudio, cursa con neovascularización coroidea originando una importante disminución en la capacidad visual.

El objetivo de este estudio era determinar la influencia de la actividad patológica (presencia de fluido retiniano) y del número total de inyecciones en la eficacia del tratamiento, en una población previamente tratada con agentes anti-VEGF.

Para ello, 33 pacientes con DMAE exudativa fueron evaluados. Se estudió la calidad visual mediante las pruebas de agudeza visual, sensibilidad al contraste y visión del color, y la integridad neuroretiniana mediante OCT. Dicha evaluación se realizó en dos visitas (PRE y POST), con una inyección anti-VEGF de diferencia entre ambas.

Los resultados PRE y POST mostraron diferencias significativas para grosor retiniano y actividad patológica. Las retinas sin actividad previa presentaron valores más moderados en el resto de las variables. Sin embargo, el tratamiento pareció ser más efectivo en pacientes con PRE-actividad, ya que la reabsorción del fluido retiniano y la consecuente disminución del grosor favorecen la mejora en la calidad visual. En cuanto al número total de inyecciones, los resultados determinaron que mayor número de inyecciones no comportaba mejoras significativas.

Como conclusión, el tratamiento para DMAE exudativa en pacientes con actividad patológica parece ser más eficaz que en aquellos sin fluido retiniano. Además, el tratamiento mantiene la calidad visual y evita la progresión de la patología más que recuperar la funcionalidad visual.



MÀSTER UNIVERSITARI EN OPTOMETRIA I CIÈNCIES DE LA VISIÓ

CARACTERISTIQUES VISUALS I MORFOLÒGIQUES EN UNA POBLACIÓ AMB DEGENERACIÓ MACULAR ASSOCIADA A L'EDAT EXUDATIVA PRE-TRACTADA

RESUM


La degeneració macular associada a l'edat (DMAE) és la principal causa de ceguesa legal en països desenvolupats. La DMAE exudativa, en la qual se centra aquest estudi, cursa amb neovascularització coroïdal originant una important disminució en la capacitat visual.

L'objectiu d'aquest estudi va ser determinar la influència de l'activitat patològica (presència de líquid retinal) i del nombre total d'injeccions en l'eficàcia del tractament en una població prèviament tractada amb agents anti-VEGF.

Així, 33 pacients amb DMAE exudativa van ser avaluats. Es va estudiar la qualitat visual mitjançant les proves d'agudes visual, sensibilitat al contrast i visió del color, i la integritat neuroretiniana mitjançant OCT. Aquesta avaluació es va realitzar en dues visites (PRE i POST), amb una injecció anti-VEGF de diferència entre ambdues.

Els resultats PRE i POST van mostrar diferències significatives per al gruix retinal i activitat patològica. Les retines sense activitat prèvia van presentar valors més moderats a la resta de les variables. No obstant, el tractament semblava ser més efectiu en pacients amb PRE-activitat, ja que la reabsorció del fluid retinal i la conseqüent disminució del gruix afavoreixen la millora en la qualitat visual. Pel que fa al nombre total d'injeccions, els resultats van determinar que un major nombre d'injeccions no comportava millores significatives.

Com a conclusió, el tractament per DMAE exudativa en pacients amb activitat patològica sembla ser més eficaç que en aquells sense fluid retinal. A més, el tractament manté la qualitat visual i evita la progressió de la patologia més que recuperar la funcionalitat visual.



MÀSTER UNIVERSITARI EN OPTOMETRIA I CIÈNCIES DE LA VISIÓ

VISUAL AND MORPHOLOGICAL CHARACTERISTICS IN A PRE-TREATED EXUDATIVE AGE-RELATED MACULAR DEGENERATION POPULATION

ABSTRACT

Age-related macular degeneration (AMD) is the leading cause of legal blindness in developed countries. Exudative AMD, in which present study is focused, presents with choroidal neovascularization inducing to an important decrease in visual capacity.

The aim of this study was to determine the influence of disease activity (presence of retinal fluid) and the total number of injections on the treatment efficacy in a pre-treated population with anti-VEGF injections.

For this, 33 patients with exudative AMD were evaluated. Visual quality was assessed by visual acuity, contrast sensitivity and colour vision tests, and neuroretinal integrity by OCT. This evaluation was performed in two clinical visits (PRE and POST), with an anti-VEGF injection of difference between both.

PRE and POST results showed significant differences for retinal thickness and disease activity. Non-active retinas displayed more moderate values in the rest of the variables. However, treatment seemed to be more effective in patients with PRE-activity, since the retinal fluid reabsorption and the consequent decrease in the retinal thicknesses favoured the visual quality improvement. Regarding the number of total injections, the results determined that more injections did not achieve a significant improvement.

In conclusion, the treatment for exudative AMD in patients with disease activity seems to be more efficient than in those without retinal fluid. In addition, treatment maintains visual quality and avoids the progression of pathology rather than visual function recovery.



COVER LETTER

Dear Editor,

Attached you will find the paper entitled “Visual and morphological characteristics in a pre-treated exudative Age-Related Macular Degeneration population”, which we are submitting for publication in *Retina* as an “Original Research paper”.

To determine whether disease activity pattern and total number of anti-VEGF injections are determining in the effectiveness of exudative AMD treatment, visual function and morphological retinal integrity were evaluated. Neuroretinal integrity and disease activity were analysed by OCT imaging as well as visual function was evaluated by visual acuity, contrast sensitivity and colour vision tests in 33 patients with exudative AMD. Results showed that neither visual nor morphological changes were significant, only substantial improvements were found in resolution of oedema and thus reduction of retinal thickness. Furthermore, no statistically differences were found according to number of previous injections, as treatment in exudative AMD appeared to maintain visual abilities.

We state that the data submitted have not been published and are not under current considerations elsewhere, that all authors have contributed significantly in the design, interpretation of data and all are in agreement with the content of the study.

We would be very grateful for any comments or suggestions you may wish to make.

Thanking you for your kind attention, we look forward to hearing from you at your convenience.

Yours sincerely,



TITLE PAGE

Title: Visual and morphological characteristics in a pre-treated exudative Age-Related Macular Degeneration population.

Abbreviated title: Outcomes of the exudative-AMD treatment

Author's name: Suárez Vidal, Antía¹

Lluch Margarit, Sara PhD²

Merindano Encina, María Dolores PhD²

¹ Terrassa School of Optics and Optometry (FOOT). Universitat Politècnica de Catalunya (UPC)

² Optics and Optometry Department. Universitat Politècnica de Catalunya (UPC)

Author's correspondence: Antía Suárez Vidal

Email address: antia.suarez3@gmail.com

Disclosure: The authors report no conflicts of interest and have no proprietary interest in any of the materials reported in this article. This research received no specific grant from funding agency in the public, commercial, or not-for-profit sectors.

KEY WORDS AND SUMMARY STATEMENT

Key Words: Age-related macular degeneration; Anti-VEGF; Colour vision; Contrast sensitivity; Choroidal neovascularization; Exudative AMD; Optical Coherence Tomography; Retinal oedema; Retinal thickness; Visual acuity

Summary Statement: Visual and morphological traits were evaluated in patients with exudative Age-Related Macular Degeneration pre-treated with anti-VEGF injections to determine the influence of disease activity and the total number of injections in the efficiency of the treatment.

ABSTRACT

Purpose – To evaluate the visual and morphological variables of a pre-treated exudative AMD population, determining the influence of the activity pattern and the total number of anti-VEGF injections in the results.

Methods – Thirty-three patients with exudative AMD were evaluated before and one month after anti-VEGF injection treatment. Visual acuity, contrast sensitivity and colour vision were analysed to assess visual function. Retinal thickness, neuroretinal integrity and disease activity pattern were evaluated by OCT imaging. Study was divided in: comparison PRE versus POST outcomes; comparison of the results according to previous disease activity pattern; comparison of the results according to the number of previous injections.

Results– One additional anti-VEGF injection only involved statistical changes in retinal thicknesses and pathology activity. Non-active PRE retinas showed moderate values for the studied variables, although treatment was more effective in active PRE retinas with retinal oedema. No statistical differences were found in the results according to the number of total injections.

Conclusions – The resolution of oedema and, thus the reduction of retinal thicknesses, seem to be the only substantial improvement that anti-VEGF treatment provides, since neither visual nor morphological changes are statistically significant. In addition, treatment in exudative AMD appeared to maintain visual abilities and prevent disease progression

INTRODUCTION

Age-related macular degeneration (AMD) is the leading cause of legal blindness among the elderly people in developed countries, accounting for 8.7% worldwide.^{1, 2, 3} In Spain, the prevalence of AMD is 3.4%, corresponding 1.5% to dry form (also known as atrophic, which includes drusen and geographic atrophy of the retinal pigment epithelium), and 1.9% to wet or exudative form which is the more aggressive form of the pathology.²

Exudative AMD is mainly characterized by choroidal neovascularization.⁴ It is thought that hypoxia is the most important factor involved in this neovascularization, causing abnormal growth of fragile blood vessels. Furthermore, a stimulation of angiogenesis by activating transcription of the gene VEGF (Vascular endothelial grow factor) has been demonstrated both in vitro and in vivo. These facts produce dysfunction of the blood-retinal barrier, resulting in leakages into the retinal tissues.¹ Consequently, the most frequent signs of exudative AMD are due to the choroidal neovascularization leakage that causes haemorrhages, hard exudates, macular serous elevation and retinal oedemas.⁴

In fact, the exudative AMD activity is ultimately based upon the presence of macular oedema. In this sense, sub-retinal or/and intra-retinal fluids, observed by optical coherence tomography (OCT) as dark empty spaces, are decisive not only to determine the disease activity, but also the treatment and follow-up protocols.^{5, 6}

In some cases, the onset of exudative AMD can be asymptomatic, especially if the visual loss occurs in the non-dominant eye.⁷ Most important symptoms are loss of central visual, metamorphopsia (visual distortion in which straight lines appear crooked), problems on dark adaptation, inability to distinguish fine detail and central blurring.^{1, 7} Moreover, 46.5% of patients with AMD present a visual acuity $<20/400$ ² and the prevalence of severe visual loss increases with age.⁸ Risk factors include age, white race, sex (women), tobacco, cataracts and family background.

The impact of AMD on emotional well-being, activities of daily living and quality of life can be profound and appears to be directly related to the severity of the pathology. Because of it, early detection and prevention of progression are the most important public health goals.⁹ Available

therapies are not causal treatments but generally, they avoid further vision loss rather than to improve vision. Anti-angiogenic agents attempt to block various steps in the pathway of angiogenesis in choroidal neovascularization.³

Three mainly types of treatment regimen are used in exudative AMD. Monthly injections with anti-VEGF agents, independently of the disease activity, have good results although many clinic visits are required. In *pro re nata* (PRN) regimen, the patients are usually given a loading dose of 3 monthly injections and then one anti-VEGF injection every 4 weeks, only in case of active disease. In *treat-and-extend* (TAE) regimen, after 3 monthly anti-VEGF injections, the intervals between treatments increase progressively, 2 weeks at a time, as long as the disease remains non-active, that is without no signs of oedemas.^{5, 10-13}

To our knowledge, there are no studies relating the visual and morphological characteristics of exudative AMD patients according to the disease activity and the extension of the treatment received.

The aim of the present study was to examine the morphological and visual variables of a pre-treated exudative AMD population, evaluating the influence of the activity pattern and the number of anti-VEGF injections in the results.

METHODS

Subjects

Thirty-three patients, previously treated for exudative age-related macular degeneration (AMD) with intravitreal injections of anti-VEGF agents, were recruited from Hospital de Terrassa- Consorci Sanitari de Terrassa (Barcelona, Spain). Inclusion criteria were: patients aged > 55 years with diagnosed exudative AMD and visual acuity (VA) of at least 49 letters on the ETDRS chart. Exclusion criteria were: presence of another ocular pathology which could alter contrast sensitivity, colour vision or visual acuity (cataract, glaucoma, high myopia, and previous retinal surgery); some cognitive disease like Alzheimer or Parkinson; and being under some treatment which could alter colour vision. All the tests were carried out only in the eye diagnosed of AMD and, in case both eyes were affected, the eye with best VA was selected. All subjects provided

written informed consent to participate in this study, which has been approved by the ethical committee of the Terrassa Health Consortium and followed the tenets of the Declaration of Helsinki.

Procedures

Visual function tests were performed with best corrected refractive error previously established by autorefraction and subjective refraction. Visual acuity was examined with Early Treatment Retinopathy Study (ETDRS) chart, which consisted of 5 letters per line in 0.1 LogMAR steps. The distance between participants and ETDRS chart was 4 meters and test was done under mesopic conditions. If participants could see at least 20 letters at a distance of 4 meters, 30 letters were added to calculate the total number score; if not, chart was then situated at 1 meter and the total number of letters seen was the final score. In this way, more precise results are obtained since it indicates the exact number of letters instead of the number of lines seen by the patients, as other optotypes do. Higher letter scores correspond to better VA. The total numbers of letters were then translated into LogMAR notation for statistical analysis.^{14, 15}

Contrast sensitivity (CS) was measured with Pelli-Robson chart, comprising 8 lines with 2 triplets (three letters) per line, 16 triplets in total. Each triplet has different contrast, which decreases line to line from 100% to 0.6%, in units of 0.15 log. Distance between patients and chart was about 1 meter, which corresponds to a spatial frequency of 1 cycle per degree. Contrast sensitivity was measured under photopic conditions (under bright-ambient room illumination). Test final score corresponds to the triplet in which at least 2 letters are correctly identified. The results were then converted into logarithmic contrast sensitivity (logCS) notation for statistical analysis. Mean normal value in healthy patients aged > 60 is 1.71 logCS.^{16, 17}

Farnsworth Dichotomous test or D-15 panel was used to examine the colour vision deficiency. This test consists in 15 caps of different colours, which participants must order according to their chromatic similarity. It was performed with a near optic correction and under photopic conditions. Results were reproduced in a diagrammatic representation in which the colour vision defect type and severity can be observed.¹⁸ Furthermore, Total Colour Difference Score (TCDS) and Confusion Index (C-index), were calculated as colour vision indexes. TCDS is a quantitative

indication of the cap transpositions, with an ideal mean value of 125.1 for the age group over 60 years; ¹⁸ CI estimates the severity of the color vision alteration, ranging from 1 up to 4 for strong vision deficiencies. ¹⁹

Central retina imaging was performed with Optovue OCT (Avanti, Optovue, USA; www.optovue.com), a system that uses 840nm wavelength light and provides 70000 A-scan/second, with an optical resolution of 5µm (axial) and 15µm (transversal). The obtained images were used to measure central retinal (CRT) and perifoveal thicknesses (PFT), considering the mean values of 256µm and 284µm respectively ²⁰ and to study some morphological traits related with exudative AMD. In that sense, the activity pattern of the pathology (non-active and active) according to the absence or the presence of retinal fluid (RF), the integrity of the ellipsoid zone (EZ, previously referred as IS/OS), ²¹ the external limiting membrane (ELM) for the outer retinal layers, and the disorganization of the retinal inner layers (DRIL) ²² were evaluated.

All tests were realized twice, before the patients were treated by an anti-VEGF intravitreal injection and one month after the treatment. Thus, results were grouped in before and after-treatment outcomes (PRE and POST groups, respectively)

According to the aims anteriorly described, this study was divided in three phases: comparison between the PRE and POST outcomes; comparison of these two groups depending on the activity pattern prior to treatment; and comparison of the POST outcomes in function of the number of injections received before to the current study.

Statistical Analysis

Statistical analysis was performed using SPSS software V23.0 (SPSS, Chicago, IL, USA). After a preliminary exploratory analysis, the quantitative variables were examined for normality using the Kolmogorov–Smirnov test/Shapiro-Wilk test, and Mann-Whitney U test or Wilcoxon signed-rank test were carried out for comparisons between groups. For categorical variables, intergroup differences were established by Chi-squared (or Fisher exact test when $n < 5$) and McNemar tests. A p value of 0.05 or less was considered to denote statistical significance throughout the study.

RESULTS

Thirty-three patients, 16 women (48.49%) and 17 men (51.52%), aged between 64 to 92 years (mean=80.78, SD=6.80) underwent the PRE-treatment testing, while only twenty-eight patients could be evaluated in the POST-treatment tests, because 5 patients did not return for the second evaluation.

1. Comparison between PRE and POST outcomes

Summary statistics and the frequency distribution of visual and morphological variables for both PRE and POST groups are shown in table 1, 2 and 3.

	PRE (n=33)				POST (n=28)			
	LogMAR	LogCS	TCDS	CI	LogMAR	LogCS	TCDS	CI
Mean	0.31	1.29	231.77	2.54	0.27	1.30	214.85	2.35
Median	0.30	1.35	236.30	2.70	0.24	1.35	221.80	2.36
SD	0.20	0.19	58.97	0.63	0.25	0.20	57.69	0.72
Min	0.00	1.05	122.70	1.13	-0.8	0.90	117.00	1.00
Max	0.70	1.65	406.80	3.54	1.10	1.62	341.90	3.67

Table 1. Descriptive statistics of visual variables for both PRE and POST groups. CI: confusion index; LogCS: contrast sensitivity logarithm; LogMAR: Logarithm of the Minimum Angle of Resolution; TCDS: Total Colour Difference Score.

	PRE (n=33)		POST (n=28)	
	CRT	PFT	CRT	PFT
Mean	274.12	270.70	240.65	255.92
Median	263.00	261.00	242.00	255.00
SD	63.56	35.95	29.71	25.71
Min	198.00	198.00	165.00	207.00
Max	448.00	362.00	309.00	320.00

Table 2. Descriptive statistics of retinal thickness for both PRE and POST groups. CRT: central retinal thickness; PFT: Perifoveal thickness, both expressed in μm

		PRE n=33	POST n=28
		%	%
ACTIVITY	Non-active	42.4	85.2
	Active	57.6	14.8
RF	Intraretinal	31.6	0.0
	Subretinal	47.4	100.0
EZ	Intra + sub	21.0	0.0
	Normal	45.5	70.4
ELM	Altered	54.5	29.6
	Normal	72.7	85.2
DRIL	Altered	27.3	14.8
	Absence	81.8	88.9
	Presence	18.2	11.1

Table 3. Frequency distribution of qualitative OCT morphological variables for both PRE and POST groups. DRIL: Disorganization of the retinal inner layers; ELM: external limiting membrane; EZ: ellipsoid zone, RF: retinal fluid.

While VA and CS values were very similar in both groups, colour indexes and some morphological variables showed a slight improvement. In that sense, CRT and PFT decreased in the second visit and the cases with active exudative AMD were reduced from 19 (57.6%) of PRE to only 4 (14.8%) of POST group. In the majority of these cases, retinal fluid showed a subretinal location (47.4% in PRE and 100% in POST). The retinal inner layers were disorganized (DRIL) in 6 PRE (18.2%) and in 3 POST (11.1%) group, while the number of patients with the outer retinal layers altered was greater either in PRE and POST group, especially EZ (Figure1).

Statistical comparison for dependent samples (Wilcoxon signed-rank and McNemar tests) displayed no significant differences ($p > 0.05$) between the visual variables of PRE and POST outcomes. This means that visual acuity, contrast sensitivity nor colour vision did not improved after anti-VEGF injection. Related to the morphological results, the variables that had a significant improvement between PRE and POST groups were CRT ($p = 0.001$), PFT ($p = 0.004$) and the activity pattern of the pathology ($p = 0.000$).

Therefore, the next phase of the present study was to analyse if the activity pattern of the disease before anti-VEGF treatment was related to the visual and morphological variables of PRE and POST groups.




Figure 1. Representative OCT images of two exudative AMD patient. A: retina with relatively preserved morphological traits, and normal central and perifoveal thickness. B: retina with affected morphological traits, and increased central and perifoveal thickness. DRIL: Disorganization of the retinal inner layers; ELM: External limiting membrane; EZ: ellipsoid zone.

2. Non-active vs active exudative AMD outcomes

According to the activity pattern before treatment, the 33 initial patients were distributed into two groups, 14 patients (42.4%) in the non-active group, and 19 patients (57.6%) in the active group. A diagram specifying the different comparisons realised in this phase can be seen in figure 2.




Figure 2. Comparisons between different groups for the second phase of the study.

2.1 Comparison between PRE groups

Summary statistics of visual variables in PRE non-active and PRE active groups are shown in Table 4A (in Supplementary material). In general, non-active retinas displayed better outcomes for VA, CS and color vision indexes than active cases, although no significant differences ($p > 0,05$) were found between both groups.

Descriptive statistics and the frequency distribution of morphological variables are shown in Table 4B and 4C respectively (in Supplementary material). CRT and PFT means were lower in non-active group, including the maximum values. Moreover, outer and inner retinal layers were more preserved in the former group. In that sense, EZ and ELM were apparently normal in 57.1% and 85.7% respectively in non-active cases, in front of 36.8% and 63.2% in active ones, being EZ more altered than ELM, in both groups. Finally, DRIL was only observed in 6 retinas (36.6%) with active exudative AMD (five of them with intraretinal fluid; Figure 3).

The Mann-Whitney U and chi-squared tests for independent samples indicated that, despite values were milder in non-active group, differences were only statistically significant for CRT ($U = 71.50$; $p = 0,025$) and DRIL ($p = 0.027$).




Figure 3. Absence/presence of DRIL according to the disease activity pattern

2.2 Comparison between POST groups

As in the previous section, no statistical differences were found for any of the studied visual variables (Table 5A; in Supplementary material). Regards to morphological variables, retinal thicknesses (Table 5B, in Supplementary material) were lower in the group that showed activity prior to the treatment, while both inner and outer retinal layers were better preserved in non-active group patients (Table 5C in Supplementary material). DRIL only appeared in 3 cases (18.8%) of the active group. Nevertheless, no statistical differences were shown with Mann-Whitney U test for quantitative variables, neither with Fisher exact test for qualitative variables.

2.3 Comparison between PRE and POST groups

To examine if previous activity pattern determined the effectiveness of treatment with anti-VEGF injection (Table 6, in Supplementary material), Wilcoxon signed-rank and McNemar tests for dependent samples were carried out. Despite the increasing number of cases with normal-

appeared retinal layers, no statistical differences were found between PRE and POST outcomes in non-active exudative AMD.

However, in the active cases, CI ($p=0.034$), CRT and PFT ($p=0.001$ and $p=0.012$ respectively; Figure 4) showed statistical improvement after treatment. The resolution of the retinal layer integrity was evident but the small number of cases for any of these variables did not allow the use of any statistical tests. Moreover, the number of retinas that continued active after the treatment was reduced in a drastic manner, remaining only 4 cases with subretinal edema.




Figure 4. Box-plot of retinal thicknesses outcomes, according to the disease activity pattern

It should be noted that although there were not significant differences between the visual variables of PRE and POST outcomes, these differences were larger in the active cases than in the no-active ones.

3. Comparison of the POST outcomes according to the total number of injections received

Since patients under study had previously undergone various treatments with anti-VEGF intravitreal injections, they were classified in two groups, according to the total number of injections received, regardless of whether they had disease activity or not. Group 1 included patients with at least 4 injections (n=13) and Group 2 patients with more than 4 injections (n=15). In this third phase of the study, only POST outcomes of both groups were compared.

Descriptive statistics and the frequency distributions of visual and morphological variables for both groups are shown in Table 7 (in Supplementary material). When visual variables were compared, the U Mann-Whitney test determined no statistical differences ($p > 0.05$) between groups, which means that visual acuity, contrast sensibility and colour vision did not improve after four treatments. With regard to retinal thicknesses, the same test indicated statistical differences in PFT ($p = 0.015$; Figure 5) but not for CRT ($p > 0.05$).




Figure 5. Box-plot of retinal thicknesses outcomes, according to the total number of injections received

In general, the morphological variables were more preserved in Group 1, showing more cases with no altered retinal layers than Group 2 (Table 7C, see annex). Only 3 cases (11.1%) in group

2, showed disorganization of the retinal inner layers, but no case was reported in group 1. While ELM and EZ were only altered in 9.1% of group 1 cases, the percentage increased to 18.8% and 43.8% respectively for group 2. Nonetheless, no statistical differences for any of the morphological variables ($p>0.05$) were found between the two groups, so the number of injections did not seem to have influenced the retinal morphology.

DISCUSSION

The purpose of this study was to evaluate visual and morphological traits of exudative AMD patients from Hospital de Terrassa-Consorci Sanitari de Terrassa (Barcelona, Spain), as well as determine the influence of the activity pattern and total anti-VEGF injections in the effectiveness of treatment.

In the first phase of the present study, the PRE and POST-treatment outcomes were compared. The attained results indicate that one additional anti-VEGF injection had not produced any statistical improvement in visual acuity, contrast sensitivity or colour vision nor in the retinal layer morphology. In contrast, retinal thickness (both CRT and PFT) and the disease activity pattern displayed statistical changes between the first and the second visit.

In fact, these variables have to be related in the sense that the resolution of oedema and, therefore, the loss of the activity by the effect of anti-VEGF injection, involved the reduction of the retinal thickness. It has to be commented, however, that mean CRT value in group 2 POST was lower than values reported after the resolution of oedemas in other maculopathies.^{22, 23} Even, it was lower than the $261.31\pm17.67\mu\text{m}$ found by Solé et al.²⁴ in 100 healthy eyes. Nevertheless, the same authors indicate that values of retinal thickness depend on the OCT instrument, given $256\mu\text{m}$ as a normal value for Optovue OCT device. It would be interesting to determine if the application of repeated injections is the cause of the maintenance of the retinal thickness below the normal values, thus affecting the visual function.

The disease activity between PRE and POST groups decreased noticeably, from 19 to only 4 patients with retinal fluid respectively. Despite this drastic oedema resolution, no substantial improvement in visual or morphological variables was observed.

These results coincide with those reported by Feigl et al ²⁵ in patients diagnosed with age-related maculopathy. The authors showed that VA, CS and colour vision, altered in the first visit study, had not improved one year later. Sabour-Pickett et al ²⁶ also showed no statistical changes in VA between baseline and after one year follow-up, but the mean foveal thickness had diminished significantly from baseline data in patients with exudative AMD. Feigl et al ²⁷ demonstrated that although retinal thickness decreased after 3 anti-VEGF injections in exudative AMD patients, VA and CS remained stable.

Regarding to morphological variables, the presence of intra or subretinal fluid modified the disposition of retinal layers, as will be discussed below.

To sum up, differences between PRE and POST outcomes would suggest the influence of the exudative AMD activity on the visual and morphological traits, and therefore its effect on the treatment effectiveness. In that sense, the analysis of how the disease activity pattern could affect the studied variables indicated that non-active PRE group exhibited better results related to VA and colour vision, as well as more preserved retinal morphology than active PRE retinas. In these active cases, the presence of oedema caused a considerable increase of retinal thickness.

As in the present study, several authors²⁸⁻³⁰ indicate that the treatment would produce the reduction of oedema and thus retinal thickness decreased more than in retinas without oedema. In that sense, Barteselli G et al ²⁸ concluded that the resolution of diabetic macular edema induced a significant decrease in the central thickness of the retina. However, unlike what happens in exudative AMD, the edema resolution was not associated with visual function recovery, as it has been proved in other macular pathologies. Yuzbasioglu et al ²⁹ showed that in patients with retinitis pigmentosa treated with anti-VEGF injections, cystoid macular edema disappeared at the same time that central macular thickness decreased and VA improved. Barone et al ³⁰ evaluated patients with pseudophakic cystoid macular edema treated with anti-VEGF injections, and concluded that as the edema resolved, VA increased and central macular thickness decreased significantly.

When comparing PRE and POST outcomes, the improvement in VA and color vision indexes, seems to be better in prior active than in non-active cases, although only CI was significant when

edema existed. In contrast, the morphological variables were more preserved in non-active group. Nevertheless, retinal thicknesses were significantly reduced in active disease group, mainly due to the disappearance of retinal oedemas that only remained in four patients with subretinal fluid. Probably, the reduction of oedemas could be also responsible for the greater proportion of retinal layers resolution in this group.

In addition, DRIL appeared mostly in cases with intraretinal fluid, and only in one case with subretinal fluid. After the treatment, the decrease in retinal thickness and the improvement of DRIL and outer retinal layers were better in active PRE group than in no-active PRE one, although in no significant way.

The inability to distinguish boundaries between inner layers (DRIL) on high-resolution OCT imaging may suggest destruction or disorganization of some axons and nuclei of amacrine, bipolar, and/or horizontal cells located in these areas.²² Pelosini et al³¹ and Grewal et al³² reported that large oedemas can snap bipolar axons and cause loss of visual signaling from photoreceptors to ganglion cells and this fact would explain why the active PRE group presented disorganised retinal inner layers as well as reduced results in visual variables.

In the same way, Radwan et al²³ demonstrated that in patients with diabetic oedema, significant correlation was found between DRIL and oedema, and that oedema resolution involved DRIL disappearance and the visual acuity improvement. Those results are in agreement with the present study, in the sense that only patients with retinal fluid (active group) presented altered retinal inner layers (DRIL) and worse visual and morphological outcomes than those patients with no edemas (non-active group). Likewise, Sun et al²² also described that DRIL was a stronger marker of VA in patients with diabetic macular edema, showing that extended DRIL implied poorer VA.

Several studies about photoreceptors restoration in different macular pathologies had indicated that EZ and ELM integrity were significantly correlated at all pre and post treatment times.³³ Moreover, the restoration of ELM was early than that of EZ and may reflect morphological and functional recovery of foveal photoreceptors. For that reason, the integrity of the external limiting membrane (ELM) rather than EZ was a better functional indicator.³⁴ In the current study, there

were always more retinas with affected EZ than ELM layer, corroborating that absence or presence of reflectivity in the EZ layer was not a good predictor for presence of photoreceptors.³⁵

In any case, the retina cell destruction may not be completely reversible after resolution of exudative DME activity, and potentially there will be some eyes in which visual function does not recover.²²

The improvement of the morphological characteristics after the treatment with anti-VEGF agents caused that the number of cases with altered morphological variables, especially in the active group, did not allow the application of comparing statistical tests. Therefore, it would be convenient to repeat this analysis with a larger number of patients.

Finally, in patients with more than 4 intravitreal injections of anti-VEGF agents, neither the visual conditions nor the morphological characteristics would improve with successive treatments. Therefore, after third or fourth injections, the effect of successive injections is basically focused on the reabsorption of the existing retinal oedema and on maintaining the visual and morphological values. In that sense, Ikuno et al ³⁶ and Zhang et al ³⁷ reported that after two and three months (respectively) with anti-VEGF treatment in patients with myopic choroidal exudativeization, treatment generates similar VA outcomes, that is, VA maintenance but no improvement.

In the current study, VA, CS and colour vision were maintained after four anti-VEGF injections. Thus, consecutive treatment seems to be a therapy to stabilize or prevent the progression of pathology rather than to solve the problems caused by the disease. Similarly, no changes were found in morphological retinal variables between both groups. Thus, retinal layers alterations caused by exudative AMD pathogenesis, were not resolved despite the number of injections applied. Therefore, the same as visual variables, the results of the present study suggest that the administration of more than four anti-VEGF injections seems to have a stabilising purpose more than a curative one.

In the same way, Sabour-Pickett et al ²⁶ reported that the greatest visual improvements in patients with exudative AMD treated with intravitreal injections are mainly achieved in the first three months of treatment. Courtney et al ³⁸ also indicated that first three months of anti-VEGF

injections in patients with AMD is a common time point where therapy may be discontinued or treatment interval may be extended, because it is when most important visual and morphological improvements occur. After this early treatment, there is currently no clear consensus regarding the optimal treatment regimen for exudative AMD patients. Recent studies have shown that the TAE regimen results in similar visual acuity outcomes as monthly injections, and that monthly injections and a TAE regimen give better visual acuity than the PRN regimen.¹⁰ Thus, PRN regimen is personalized treatment but it presents the disadvantage that the lesion is retreat when it has already reactivated. In contrast, monthly and TAE protocols are based on retreatment before lesion reactivating but with different number of clinic visits.^{5, 6, 10, 11, 13}

As a conclusion, the resolution of oedema and, thus the reduction of retinal thicknesses, seem to be the only substantial improvement that anti-VEGF treatment provides, since neither visual nor morphological changes are statistically significant. However, patients often perceive some visual enhancements after treatment. It could be possible that, although no significant, the small achieved changes in VA, CS and color vision may be important enough to really represent an improvement in the visual perception of these patients. Further studies with a larger population and longer period of follow-up would be of interest to corroborate this new perspective.

Acknowledgements: The authors thank to Laura Gutiérrez, ophthalmologist from Hospital de Terrassa-Consorci Sanitari de Terrassa (CST), for medical assistance and for providing patients.

REFERENCES

1. Christoforidis JB, Tecce N, Dell'Omo R. et al. Age related macular degeneration and visual disability. *Curr Drug Targets*. 2011; 12(2): 221-33.
2. Spanish Eyes Epidemiological (SEE) Study Group. Prevalence of age-related macular degeneration in Spain. *Br J Ophthalmol*. 2011; 95: 335-40.
3. Nowak JZ. Age-related macular degeneration (AMD): pathogenesis and therapy. *Pharmacol Rep*. 2006; 58(3): 353-63.
4. Kanski JJ. Capítulo 13: Trastornos maculares adquiridos. *Oftalmología clínica*. 5º Ed. UK: ELSEVIER.
5. Arnold JJ, Markey CM, Kurstjens NP, et al. The role of sub-retinal fluid in determining treatment outcomes in patients with neovascular age-related macular degeneration – a phase IV randomised clinical trial with ranibizumab: the FLUID study. *BMC Ophthalmol*. 2016; 16: 31.
6. Arias Barquet L, Monés J. New treatment protocols and follow-up in patients with exudative age-related macular degeneration. *Arch Soc Esp Oftalmol*. 2012; 87(1): 10.
7. Hogg RE, Chakravarthy U. Visual function and dysfunction in early and late age-related maculopathy. *Prog Retin Eye Res*. 2006; 25(3): 249-76.
8. Sjöstrand J, Laatikainen L, Hirvelä H, et al. The decline in visual acuity in elderly people with healthy eyes or eyes with early age-related maculopathy in two Scandinavian population samples. *Acta Ophthalmol*. 2011; 89(2): 116-23.
9. Lovie-Kitchin J, Feigl B. Assessment of age-related maculopathy using subjective vision tests. *Clin Exp Optom*. 2005; 88(5): 292-303.
10. Kvannli L, Krohn J. Switching from pro re nata to treat-and-extend regimen improves visual acuity in patients with neovascular age-related macular degeneration. *Acta Ophthalmol*. 2017.

11. Vardarinos A, Gupta N, Janjua R, et al. 24-month clinical outcomes of a treat-and-extend regimen with ranibizumab for wet age-related macular degeneration in a real life setting. *BMC Ophthalmol.* 2017; 17(1): 58.
12. Holz FG, Amoaku W, Donate J, et al. Safety and efficacy of a flexible dosing regimen of ranibizumab in neovascular age-related macular degeneration: the SUSTAIN study. *Ophthalmology.* 2011; 118(4): 663-71.
13. Berg K, Pedersen TR, Sandvik L, et al. Comparison of ranibizumab and bevacizumab for neovascular age-related macular degeneration according to LUCAS treat-and-extend protocol. *Ophthalmology.* 2015; 122(1): 146-52.
14. Peter K. Kaiser. Prospective evaluation of visual acuity assesment: a comparison of Snellen versus ETDRS charts in clinical practice (An AOS Thesis). *Trans Am Ophthalmol Soc.* 2009; 107: 311-24.
15. Vecilla M. Capítulo 1: Agudeza visual. *Manual de Optometría.* Madrid, Editorial Médica Panamericana, 2010. 3-21
16. Mäntyjärvi M, Laitinen T. Normal values for the Pelli-Robson contrast sensitivity test. *J Cataract Refract Surg.* 2001; 27(2): 261-6.
17. Maynard ML, Zele AJ, Feigl B. Mesopic Pelli-Robson contrast sensitivity and MP-1 microperimetry in healthy ageing and age-related macular degeneration. *Acta Ophthalmol.* 2016; 94(8): 772-778.
18. Bowman KJ. A method for quantitative scoring of the Farnsworth Panel D-15. *Acta Ophthalmol.* 1982; 60(6): 907-16.
19. Vingrys AJ, King-Smith PE. A quantitative scoring technique for panel tests of color vision. *Invest Ophthalmol Vis Sci.* 1988; 29(1): 50-63.
20. Solé González L, Abreu González R, Alonso Plasencia M, et al. Normal volume and thickness in spectral-domain OCT. Literature revision. Available on: <http://sociedadcanariadeoftalmologia.com/wp-content/revista/revista-23/23sco08.htm>

21. Staurenghi G, Sadda S, Chakravarthy U, et al. Proposed lexicon for anatomic landmarks in normal posterior segment spectral-domain optical coherence tomography: the IN-OCT consensus. *Ophthalmology*. 2014; 121(8) 1572-8.
22. Sun JK, Radwan SH, Soliman AZ, et al. Neural Retinal Disorganization as a Robust Marker of Visual Acuity in Current and Resolved Diabetic Macular Edema. *Diabetes*. 2015; 64(7): 2560-70.
23. Radwan SH, Soliman AZ, Tokarev J, et al. Association of Disorganization of Retinal Inner Layers With Vision After Resolution of Center-Involved Diabetic Macular Edema. *JAMA Ophthalmol*. 2015; 133(7): 820-5.
24. Solé González L, Abreu González R, Alonso Plasencia M, et al. Normal macular thickness and volume using spectral domain optical coherence tomography in a reference population. *Arch Soc Esp Oftalmol*. 2013; 88 (9): 352-8.
25. Feigl B, Brown B, Lovie-Kitchin J. et al. Monitoring retinal function in early age-related maculopathy: visual performance after 1 year. *Eye (Lond)* 2005; 19 (11): 1169-77.
26. Sabour-Pickett S, Loughman J, Nolan JM. et al. Visual performance in patients with neovascular age-related macular degeneration undergoing treatment with intravitreal ranibizumab. *J Ophthalmol*. 2013.
27. Feigl B, Greaves A, Brown B. Fuctional outcomes after multiple treatments with ranibizumab in neovascular age-related macular degeneration beyond visual acuity. *Clin Ophthalmol*. 2007; 1(2): 167-75.
28. Barteselli G, Kozak I, El-Emam S, et al. 12-month results of the standardised combination therapy for diabetic macular oedema: intravitreal bevacizumab and navigated retinal photocoagulation. *Br J Ophthalmol*. 2014; 98(8): 1036-41.
29. Yuzbasioglu E, Artunay O, Rasier R, et al. Intravitreal bevacizumab (Avastin) injection in retinitis pigmentosa. *Curr Eye Res*. 2009; 34(3): 231-7.
30. Barone A, Russo V, Prascina F, et al. Short-term safety and efficacy of intravitreal bevacizumab for pseudophakic cystoid macular edema. *Retina*. 2009; 29(1): 33-7.

31. Pelosini L, Hull CC, Boyce JF, et al. Optical coherence tomography may be used to predict visual acuity in patients with macular edema. *Invest Ophthalmol Vis Sci.* 2011; 52(5): 2741-8.
32. Grewal DS, Hariprasad SM, Jaffe GJ. Role of Disorganization of Retinal Inner Layers as an Optical Coherence Tomography Biomarker in Diabetic and Uveitic Macular Edema. *Ophthalmic Surg Lasers Imaging Retina.* 2017; 48(4): 282-288.
33. Mitamura Y, Mitamura-Aizawa S, Katome T, et al. Photoreceptor impairment and restoration on optical coherence tomographic image. *J Ophthalmol.* Volume 2013, Article ID 518170, 7 pages
34. Wakabayashi T, Fujiwara M, Sakaguchi H, et al. Foveal microstructure and visual acuity in surgically closed macular holes: spectral-domain optical coherence tomographic analysis. *Ophthalmology.* 2010; 117(9): 1815-24.
35. Scoles D, Flatter JA, Cooper RF, et al. Assessing photoreceptor structure associated with ellipsoid zone disruptions visualized with optical coherence tomography. *Retina.* 2016; 36(1): 91-103.
36. Ikuno Y, Ohno-Matsui K, Wong TY, et al. Intravitreal Aflibercept Injection in Patients with Myopic Choroidal Neovascularization: The MYRROR Study. *Ophthalmology.* 2015; 122(6): 1220-7.
37. Zhan Y, Han Q, Ru Y, et al. Anti-VEGF treatment for myopic choroid neovascularization: from molecular characterization to update on clinical application. *Drug Des Devel Ther.* 2015; 9: 3413-21.
38. Courtney RJ, McClintic JJ, Ehlers JP. Comparison of spectral domain optical coherence tomography scan patterns and clinical review strategies in neovascular age-related macular degeneration. *Retina.* 2015; 35(7): 1315-22.

SUPPLEMENTARY MATERIAL

A	NON-ACTIVE eAMD (n=14)				ACTIVE eAMD (n=19)			
	LogMAR	LogCS	TCDS	CI	LogMAR	LogCS	TCDS	CI
Mean	0.28	1.23	213.76	2.35	0.33	1.33	245.05	2.67
Median	0.23	1.20	222.85	2.56	0.34	1.35	246.60	2.81
SD	0.22	0.20	53.10	0.71	0.19	0.18	60.89	0.54
Min	0.00	1.05	122.70	1.13	0.06	1.05	163.90	1.60
Max	0.66	1.65	303.40	3.38	0.70	1.65	406.80	3.54

B	NON-ACTIVE eAMD (n=14)		ACTIVE eAMD (n=19)	
	CRT	PFT	CRT	PFT
Mean	245.29	260.07	295.37	278.53
Median	237.50	253.50	269.00	270.00
SD	30.78	25.09	72.85	41.11
Min	198.00	226.00	208.00	198.00
Max	316.00	322.00	448.00	362.00

C	NON-ACTIVE eAMD (n=14)		ACTIVE eAMD (n=19)	
		%		%
EZ	Normal	57.1		36.8
	Altered	42.9		63.2
ELM	Normal	85.7		63.2
	Altered	14.3		36.8
DRIL	Absence	100.0		68.4
	Presence	0.0		31.6

Table 4. PRE-group outcomes, according to the activity pattern prior to treatment. A: descriptive statistics for the studied visual variables; B: descriptive statistics for retinal thicknesses; C: frequency distribution of the studied morphological variables. CI: confusion index; CRT: central retinal thickness; DRIL: Disorganization of the retinal inner layers; ELM: external limiting membrane; EZ: ellipsoid zone; LogCS: contrast sensitivity logarithm; LogMAR: Logarithm of the Minimum Angle of Resolution; PFT: Perifoveal thickness; TCDS: Total Colour Difference Score. Thickness expressed in μm .

A	NON-ACTIVE eAMD (n=12)				ACTIVE eAMD (n=16)			
	LogMAR	LogCS	TCDS	CI	LogMAR	LogCS	TCDS	CI
Mean	0.21	1.32	215.18	2.34	0.27	1.32	220.58	2.43
Median	0.20	1.35	199.40	2.11	0.30	1.35	239.60	2.72
SD	0.187	0.20	71.01	0.90	1.98	0.20	49.10	0.60
Min	-0.06	1.05	117.00	1.00	-0.08	0.90	117.00	1.00
Max	0.64	1.65	341.90	3.67	0.60	1.65	281.50	3.10

B	NON-ACTIVE eAMD (n=12)		ACTIVE eAMD (n=16)	
	CRT	PFT	CRT	PFT
Mean	243.00	261.36	238.93	251.93
Median	243.00	254.00	241.00	256.00
SD	29.53	28.17	30.76	23.93
Min	191.00	226.00	165.00	207.00
Max	309.00	320.00	302.00	304.00

C	NON-ACTIVE eAMD (n=11)		ACTIVE eAMD (n=16)	
		%		%
EZ	Normal	81.8		81.3
	Altered	18.2		18.7
ELM	Normal	90.9		62.5
	Altered	9.1		37.5
DRIL	Absence	100.0		81.3
	Presence	0.0		18.7

Table 5. POST-group outcomes, according to the activity pattern prior to treatment. A: descriptive statistics for the studied visual variables; B: descriptive statistics for retinal thicknesses; C: frequency distribution of the studied morphological variables. CI: confusion index; CRT: central retinal thickness; DRIL: Disorganization of the retinal inner layers; ELM: external limiting membrane; EZ: ellipsoid zone; LogCS: contrast sensitivity logarithm; LogMAR: Logarithm of the Minimum Angle of Resolution; PFT: Perifoveal thickness; TCDS: Total Colour Difference Score. Thickness expressed in μm .

A	NON-ACTIVE eAMD		ACTIVE eAMD	
	PRE (n=14)	POST (n=11)	PRE (n=19)	POST (n=16)
LogMAR	0.28 (0.22)	0.21 (0.19)	0.33 (0.19)	0.27 (1.98)
LogCS	1.23 (0.20)	1.32 (0.20)	1.33 (0.18)	1.32 (0.20)
TCDS	213.76 (53.10)	215.18 (71.01)	245.05 (60.89)	220.58 (49.10)
CI	2.35 (0.71)	2.34 (0.90)	2.67 (0.54)	2.43 (0.60)

B	NON-ACTIVE eAMD		ACTIVE eAMD	
	PRE (n=14)	POST (n=11)	PRE (n=19)	POST (n=16)
CRT	245.29 (30.78)	243.00 (29.53)	295.37 (72.85)	238.93 (30.76)
PFT	260.07 (25.09)	261.36 (28.17)	278.53 (41.11)	251.93 (23.93)

C		% NON-ACTIVE eAMD		% ACTIVE eAMD	
		PRE (n=14)	POST (n=11)	PRE (n=19)	POST (n=16)
EZ	Normal	57.1	81.8	36.8	62.5
	Altered	42.9	18.2	63.2	37.5
ELM	Normal	85.7	90.9	63.2	81.3
	Altered	14.3	9.1	36.8	18.7
DRIL	Absence	100.0	100.0	68.4	81.3
	Presence	0.0	0.0	31.6	18.7
ACTIVITY	No-active	100.0	100.0	0.0	75.0
	Active	0.0	0.0	100	25.0

Table 6. PRE and POST-group outcomes, according to the activity pattern prior to treatment. A: Mean (SD) for the studied visual variables; B: Mean (SD) for retinal thicknesses; C: frequency distribution of the studied morphological variables. CI: confusion index; CRT: central retinal thickness; DRIL: Disorganization of the retinal inner layers; ELM: external limiting membrane; EZ: ellipsoid zone; LogCS: contrast sensitivity logarithm; LogMAR: Logarithm of the Minimum Angle of Resolution; PFT: Perifoveal thickness; TCDS: Total Colour Difference Score. Thickness expressed in μm .

A	GROUP 1 (n=13)				GROUP 2 (n=15)			
	LogMAR	LogCS	TCDS	CI	LogMAR	LogCS	TCDS	CI
Mean	0.26	1.28	195.25	2.13	0.29	1.32	231.84	2.55
Median	0.18	1.35	169.50	1.96	0.30	1.35	239.60	2.72
SD	0.32	1.90	63.57	0.82	0.17	0.21	47.84	0.57
Min	-0.06	1.05	117.00	1.00	-0.08	0.90	146.80	1.41
Max	1.10	1.65	293.60	3.26	0.60	3.67	341.90	3.67

B	GROUP 1 (n=11)		GROUP 2 (n=15)	
	CRT	PFT	CRT	PFT
Mean	252.64	271.00	231.87	244.87
Median	245.00	263.00	240.00	242.00
SD	28.86	26.15	28.04	19.54
Min	223.00	233.00	165.00	207.00
Max	309.00	320.00	262.00	277.00

C	GROUP 1 (n=11)		GROUP 2 (n=16)	
		%		%
ELM	Normal	9.91		81.25
	Altered	9.09		18.75
EZ	Normal	90.91		56.30
	Altered	9.09		43.80
DRIL	Absence	100		81.25
	Presence	-		18.75

Table 7. POST-group outcomes, according to the total number of injections received: GROUP 1: ≤ 4 injections GROUP 2: > 4 injections. A: descriptive statistics for the studied visual variables; B: descriptive statistics for retinal thicknesses; C: frequency distribution of the studied morphological variables. CI: confusion index; CRT: central retinal thickness; DRIL: Disorganization of the retinal inner layers; ELM: external limiting membrane; EZ: ellipsoid zone; LogCS: contrast sensitivity logarithm; LogMAR: Logarithm of the Minimum Angle of Resolution; PFT: Perifoveal thickness; TCDS: Total Colour Difference Score. Thickness expressed in μm .

Author Resources



Author Resources

- Instructions for Authors (this page)
- [Reprint Ordering](#)
- [Permissions Requests](#)
- [Reprints](#)

Scope: RETINA®, The Journal of Retinal and Vitreous Diseases, publishes original and special articles concerning disorders of the retina and vitreous.

Ethical/Legal Considerations. A submitted manuscript must be an original contribution not previously published (except as an abstract or preliminary report), must not be under consideration for publication elsewhere, and if accepted, it must not be published elsewhere in similar form, in any language, without the consent of Lippincott Williams & Wilkins. Each person listed as an author is expected to have participated in the study to a significant extent. Although the editors and referees make every effort to ensure the validity of published manuscripts, the final responsibility rests with the authors, not with the Journal, its editors, or the publisher. **All manuscripts must be submitted on-line through the journal's Web site at www.retinajournal.com.** See submission instructions under "On-line manuscript submission."

FINANCIAL DISCLOSURE: Each author warrants that he or she has no commercial associations (e.g., consultancies, stock ownership, equity interest, patent/licensing arrangements, etc.) that might pose a conflict of interest in connection with the submitted article, except as disclosed on a separate attachment. All funding sources supporting the Work and all institutional or corporate affiliations of the authors are acknowledged in a footnote in the Work and on the Copyright/Financial Disclosure form that is requested upon acceptance.

Patient anonymity and informed consent. It is the author's responsibility to ensure that a patient's anonymity be carefully protected and to verify that any experimental investigation with human subjects reported in the manuscript was performed with informed consent and following all the guidelines for experimental investigation with human subjects required by the institution(s) with which all the authors are affiliated. Authors should mask patients' eyes and remove patients' names from figures unless they obtain written consent from the patients and submit written consent with the manuscript.

Required Author Forms: Several forms are required of all authors at the time a manuscript is submitted for publication. These include the following:

- **Authorship Responsibility, Disclosure, and Copyright Transfer:** All authors must sign a copy of the journal's "Authorship Responsibility, Financial Disclosure, and Copyright Transfer" form and submit it at the time of manuscript submission. The Financial Disclosure section of this form is required by the International Committee of Medical Journal Editors (ICMJE).

Authorship Criteria Form: Each author must also fill out the Authorship Form confirming that all persons who have contributed substantially but who are not authors are identified in the Acknowledgment section and that written permission from each person acknowledged has been obtained. The criteria of authorship are defined by the International Committee of Medical Journal Editors (ICMJE).

Questions? Contact the Editorial Office at retina@retinajournal.com.

Compliance with NIH and Other Research Funding Agency Accessibility Requirements

A number of research funding agencies now require or request authors to submit the post-print (the article after peer review and acceptance but not the final published article) to a repository that is accessible online by all without charge. As a service to our authors, LWW will identify to the National Library of Medicine (NLM) articles that require deposit and will transmit the post-print of an article based on research funded in whole or in part by the National Institutes of Health, Wellcome Trust, Howard Hughes Medical Institute, or other funding agencies to PubMed Central. The revised Copyright Transfer Agreement provides the mechanism.

Copyright. In view of the Copyright Revision Act of 1976, effective January 1, 1978, all transmittal letters must contain the following language before manuscripts can be reviewed for possible publication:

"In consideration of the journal RETINA® taking action in reviewing and editing my (our) submission, which represents an original article (the copyright of which is held solely by the undersigned), the author(s) undersigned hereby transfers, assigns, or otherwise conveys all copyright ownership to the Ophthalmic Communications Society, Inc. in the event that such work is published in the journal RETINA®. Manuscripts authored by US government employees are exempted from the requirement of the preceding paragraph if they constitute works of the US government. Under copyright law, a work of authorship prepared by an officer or employee of the US government as part of that person's official duties is in the public domain and is not subject to copyright protection. All manuscripts constituting such works that are submitted to RETINA®(or their accompanying transmittal letters) should not contain the language quoted in the preceding sentences, but must contain the following language instead, accompanied by the signature(s) of the author(s), before they can be considered for possible publication: "The submitted manuscript is a work of authorship prepared as part of my (our) official duties as an officer or employee of the US government and is therefore in the public domain."

OPEN ACCESS

LWW's hybrid open access option is offered to authors whose articles have been accepted for publication. With this choice, articles are made freely available online immediately upon publication. Authors may take advantage of the open access option at the point of acceptance to ensure that this choice has no influence on the peer review and acceptance process. These articles are subject to the journal's standard peer-review process and will be accepted or rejected based on their own merit.

Authors of accepted peer-reviewed articles have the choice to pay a fee to allow perpetual unrestricted online access to their published article to readers globally, immediately upon publication. The article processing charge for Retina, the *Journal of Retinal and Vitreous Diseases*, is \$3,100. The article processing charge for authors funded by the Research Councils UK (RCUK) is \$3,900. The publication fee is charged on acceptance of the article and should be paid within 30 days by credit card by the author, funding agency or institution. Payment must be received in full for the article to be published open access. Any additional standard publication charges, such as for color images, will also apply.

- ***Authors retain copyright***

Authors retain their copyright for all articles they opt to publish open access. Authors grant LWW a license to publish the article and identify itself as the original publisher.

In addition, each author must complete and submit the journal's copyright transfer agreement, which includes a section on the disclosure of potential conflicts of interest based on the recommendations of the International Committee of Medical Journal Editors, "Uniform Requirements for Manuscripts Submitted to Biomedical Journals" (www.icmje.org/update.html).

A copy of the form is made available to the submitting author within the Editorial Manager submission process. Co-authors will automatically receive an Email with instructions on completing the form upon submission.

- ***Creative Commons license***

Articles opting for open access will be freely available to read, download and share from the time of publication. Articles are published under the terms of the Creative Commons License Attribution-Non Commercial No Derivative 3.0 which allows readers to disseminate and reuse the article, as well as share and reuse of the scientific material. It does not permit commercial exploitation or the creation of derivative works without specific permission. To view a copy of this license visit:

<http://creativecommons.org/licenses/by-nc-nd/3.0>.

- ***Compliance with NIH, RCUK, Wellcome Trust and other research funding agency accessibility requirements***

A number of research funding agencies now require or request authors to submit the post-print (the article after peer review and acceptance but not the final published article) to a repository that is accessible online by all without charge. As a service to our authors, LWW identifies to the National Library of Medicine (NLM) articles that require deposit and transmits the post-print of an article based on research funded in whole or in part by the National Institutes of Health, Howard Hughes Medical Institute, or other funding agencies to PubMed Central. The revised Copyright Transfer Agreement provides the mechanism. LWW ensures that authors can fully comply with the public access requirements of major funding bodies worldwide. Additionally, all authors who choose the open access option will have their final published article deposited into PubMed Central.

RCUK and Wellcome funded authors can choose to publish their paper as open access with the payment of an article process charge (gold route), or opt for their accepted manuscript to be deposited (green route) into PMC with an embargo.

With both the gold and green open access options, the author will continue to sign the Copyright Transfer Agreement (CTA) as it provides the mechanism for LWW to ensure that the author is fully compliant with the requirements. After signature of the CTA, the author will then sign a License to Publish where they will then own the copyright. Those authors who wish to publish their article via the gold route will be able to publish under the terms of the Attribution 3.0 (CCBY) License. To view of a copy of this license visit: <http://creativecommons.org/licenses/by/2.0/>. Those authors who wish to publish their article via the green route will be able to publish under the rights of the Attribution Non-commercial 3.0 (CCBY NC) license (<http://creativecommons.org/licenses/by-nc/2.0/>).

It is the responsibility of the author to inform the Editorial Office and/or LWW that they have RCUK funding. LWW will not be held responsible for retroactive deposits to PMC if the author has not completed the proper forms.

FAQ for open access

<http://links.lww.com/LWW-ES/A48>

Permissions. Authors must submit written permission from the copyright owner (usually the publisher) to use direct quotations, tables, or illustrations that have appeared in copyrighted form elsewhere, along with complete details about the source. Any permissions fees that might be required by the copyright owner are the responsibility of the authors requesting use of the borrowed material, not the responsibility of Lippincott Williams & Wilkins.

For permission and/or rights to use content for which the copyright holder is LWW or the society, please go to the journal's website and after clicking on the relevant article, click on the "Request Permissions" link under the "Article Tools" box that appears on the right side of the page. Alternatively, send an e-mail to customercare@copyright.com.

For Translation Rights & Licensing queries, contact Silvia Serra, Translations Rights, Licensing & Permissions Manager, Wolters Kluwer Health (Medical Research) Ltd, 250 Waterloo Road, London SE1 8RD, UK. Phone: +44 (0) 207 981 0600. E-mail: silvia.serra@wolterskluwer.com

For Special Projects **and Reprints** (U.S./Canada), contact Alan Moore, Director of Sales, Lippincott Williams & Wilkins, Two Commerce Square, 2001 Market Street, Philadelphia, PA 19103. Phone: 215-521-8638. E-mail: alan.moore@wolterskluwer.com

For Special Projects **and Reprints** (non-U.S./Canada), contact Silvia Serra, Translations Rights, Licensing & Permissions Manager, Wolters Kluwer Health (Medical Research) Ltd, 250 Waterloo Road, London SE1 8RD, UK. Phone: +44 (0) 207 981 0600. E-mail:

silvia.serra@wolterskluwer.com

Manuscript Submission

On-line manuscript submission: All manuscripts must be submitted on-line through the Web site: **www.retinajournal.com**. **First-time users:** Please click the Register button from the menu and enter the requested information. On successful registration, you will be sent an e-mail indicating your user name and password. Print a copy of this information for future reference. *Note:* If you have received an e-mail from us with an assigned user ID and password, or if you are a repeat user, do not register again. Just log in. Once you have an assigned ID and password, you do not have to re-register, even if your status changes (that is, author, reviewer, or editor).

Authors: Please click the log-in button from the menu at the top of the page and log into the system as an Author. Submit your manuscript according to the author instructions. You will be able to track the progress of your manuscript through the system. If you experience any problems, please contact us either by e-mail at **retina@retinajournal.com** or 610-525-4849.

PREPARATION OF MANUSCRIPTS

Manuscripts that do not adhere to the following instructions will be returned to the corresponding author for technical revision before undergoing peer review. The manuscript/text part of your submission must be in WORD (.DOC or .DOCX format)

Title Page. Page 1 of your manuscript. Make the full title brief but meaningful. Make it explicit for indexing. The abbreviated title cannot be greater than 40 characters. List all authors by first name, middle, initial and last name, with highest academic or medical degree. When applicable, the footnote should include each author's departmental affiliation and the institution where the study was performed, grants and funds in support of the study, and if the paper was presented at

a meeting, the name of the organization, place, and date on which it was read. The name and mailing address of the author to whom correspondence and requests for reprints should be directed must be provided. All manuscripts must have a sentence indicating whether any authors have a proprietary interest.

The title page must also include disclosure of funding received for this work from any of the following organizations: National Institutes of Health (NIH); Wellcome Trust; Howard Hughes Medical Institute (HHMI); and other(s).

Key Words and Summary Statement. Page 2 of your manuscript. List words and terms in alphabetic order under which you believe the article should be indexed. The editorial board reserves the right to alter this list as it deems necessary. You may list up to 10 key words. Also include a brief summary statement (50 words or less) to be printed in the table of contents.

Structured Abstract. Page 3 of your manuscript. A structured abstract of not more than 200 words is required. The abstract should be divided into four sections with the following headings: Purpose, Methods, Results, and Conclusion.

Text. The introduction should be pertinent to the study but not an in-depth review of the literature. The materials, subjects, and methods should be clearly defined so that the study may be duplicated by other investigators. Results should be as concise as possible. The discussion is an explanation of the results of the study and should limit itself to the subject matter of the paper.

Visual acuities are to be reported using Snellen visual acuity ratios in all clinically applicable situations. Snellen acuities may be placed in parentheses next to other forms of reporting of visual acuities such as logMAR. The use of logMAR reporting is permissible but only when used for the purpose of statistical analysis. Decimals are to be used only when Snellen ratios are also provided. The decimal system alone for reporting visual acuities is NOT acceptable for the journal RETINA. Improvements of visual acuities are to be reported in letters of improvement (ETDRS) or in lines of improvement using ETDRS letter scores. These changes should be made where applicable in the abstract, manuscript text, figure legends and all tables.

Tables and figures are to be separate files that are submitted individually and cannot be embedded in the word document

Tables. (Double space, remove) Each table must be submitted in a separate WORD (.DOC or .DOCX). Do not embed Tables in your text. Number consecutively as they are presented in the text. A brief title for each table must be supplied. Give each column a short or abbreviated heading. Place explanatory matter in the footnotes, not in the heading. The following symbols are to be used in this sequence asterisk (*), dagger (†), upside-down-dagger, double-dagger (§), double asterisks (*), 2 daggers, 2 upside-down-daggers, 2 double-daggers. Horizontal and vertical rules are to be omitted. Previously published data must be acknowledged fully and written permission for its use must be obtained. Each table should be self-explanatory, i.e., all

information necessary to understand the table without the use of the text should be contained in the caption and the table itself. Due to space considerations, do not use a table for data that can be given in the text in one or two sentences.

Figures. Figures must be submitted as individual files. They cannot be embedded in the word document. Cite figures consecutively in the text, and number them in the order in which they are discussed. If the images are not of a high enough resolution to permit quality reproduction for publication purposes, they will be returned to the author.

Digital art should be created/scanned and saved and submitted as either a TIFF (tagged image file format), an EPS (encapsulated postscript) file. **Electronic photographs—radiographs, CT scans, and so on—and scanned images must have a resolution of at least 300 dpi. Line art must have a resolution of at least 1200 dpi (dots per inch).** If fonts are used in the artwork, they must be converted to paths or outlines or they must be embedded in the files. **Color images must be created/scanned and saved and submitted as CMYK files.**

Color figures. The journal encourages the submission of color art as an important part of image recognition in ophthalmology. Color costs are to be paid by the authors. Authors will be charged at the rate of \$500 for the first figure within the article, \$100 for each additional single-image figure within the same article, or \$200 for each additional figure with more than one part (labeled "a," "b," etc.).

Detailed Figure Instructions: For a step by step guide for submitting Digital Art to please visit <http://links.lww.com/ES/A42>. Click "For Authors" then click "Artwork" in the menu to the right. Visit the "Digital Art Checklist" and "5 Steps for Creating Digital Artwork" for specific guidelines.

Figure legends. Legends must be submitted for all figures. They should be brief and specific. Legends should appear on a single page after your references. Use scale markers in the image for electron micrographs, and indicate the type of stain used.

Supplemental Digital Content

Supplemental Digital Content (SDC): Authors may submit SDC via Editorial Manager to LWW journals that enhance their article's text to be considered for online posting. SDC may include standard media such as text documents, graphs, audio, video, etc. On the Attach Files page of the submission process, please select Supplemental Audio, Video, or Data for your uploaded file as the Submission Item. If an article with SDC is accepted, our production staff will create a URL with the SDC file. The URL will be placed in the call-out within the article. SDC files are not

copy-edited by LWW staff, they will be presented digitally as submitted. For a list of all available file types and detailed instructions, please visit <http://links.lww.com/A142>.

SDC Call-outs

Supplemental Digital Content must be cited consecutively in the text of the submitted manuscript. Citations should include the type of material submitted (Audio, Figure, Table, etc.), be clearly labeled as "Supplemental Digital Content," include the sequential list number, and provide a description of the supplemental content. All descriptive text should be included in the call-out as it will not appear elsewhere in the article.

Example:

We performed many tests on the degrees of flexibility in the elbow (see Video, Supplemental Digital Content 1, which demonstrates elbow flexibility) and found our results inconclusive.

List of Supplemental Digital Content

A listing of Supplemental Digital Content must be submitted at the end of the manuscript file. Include the SDC number and file type of the Supplemental Digital Content. This text will be removed by our production staff and not be published.

Example:

Supplemental Digital Content 1. wmv

SDC File Requirements

Most acceptable file types are permissible up to 10 MBs. Audio or video files can be up to 100 MBs. For a list of all available file types and detailed instructions, please visit <http://links.lww.com/A142>.

Accessing SDC files

All acceptable file types should be accessible using your default/preferred media player (e.g., Windows Media Player, Quicktime, etc.) If you experience difficulties accessing supplemental video files, you may also consider using the free VLC media player, accessible via <http://www.videolan.org>.

Acknowledgments. Sponsoring organizations and grants should be acknowledged in a footnote on the title page. Only rarely does an exceptional contribution merit an acknowledgment in RETINA®.

References. References must be numbered consecutively, according to their appearance in text. Once a reference is cited, all subsequent citations should be to the original number. Personal communications and unpublished data should be limited and incorporated into text without a reference number. Reference to studies that have been accepted but not yet published should

indicate where they will be published. References should be cited in text as follows: Brown and Jones,¹ Smith,² and Johnson et al³ described. The term **et al** is reserved for articles with three or more authors. The author is responsible for complete and accurate references, including the proper capitalization and accent marks used in foreign-language publications.

In listed references, all names of authors should be given, unless there are more than four. If there are five or more, list the first three, followed by "et al." References to journal articles should include (1) author, (2) title, (3) journal name (as abbreviated in **Index Medicus**), (4) year, (5) volume number, and (6) inclusive page numbers, in that order. References to books should include (1) author(s), (2) chapter title (if any), (3) editor (if any), (4) title of book, (5) city of publication, (6) publisher, (7) year, and (8) page numbers. Volume and edition numbers, and names of translator should be included when appropriate.

The following style is used by the Journal for periodicals (1) and for books (2):

1. Smith BR. Diseases of the retina. Retina 1981; 1:25-28.
2. Jones IS, Jakobiec FA. Diseases of the Orbit. Hagerstown, MD: Harper & Row, 1979;9-17.

If there is any doubt about abbreviation of a journal name, it should be spelled out completely.

Letters to the Editor:

Since 2014, all Letters to the Editor are published online-only and within the iPad App. There are different types of Letters to the Editor (LTE). If the purposes of the LTE is to comment on a published article, the first sentence of the LTE should include the name of the published article's first author along with the title of the published article. The article discussed should then be referenced at this time or should be referenced at the end of the LTE. If the LTE is a reply to a previously submitted LTE, the first sentence should include the name of the letter's author and cite the letter as reference. The previously published article should then be referenced as well either in the body of the text or at the end of the response to the LTE. If the LTE is a reply/response which was solicited by the editors of RETINA, please submit the reply through the "Invited Commentary" process. Do not submit it as an original article or submission. Signed copyright transfer forms and Authorship forms from each author is required with your submission.

Other Sections. The journal also accepts manuscripts for other sections such as Review Articles, Diagnostic & Therapeutic Challenges, Correspondences (Letter to the Editor, and Response to Letter to the Editor), Clinicopathologic Correlations, Surgical Techniques, and New Instruments. All manuscripts must be double-spaced, include a Title Page and be no more than 5 pages of manuscript text in length, and should include no more than 4 figures and 5 references. The format for all submitted manuscripts is basically as described above with a few exceptions. Review Articles are not limited in length. Diagnostic & Therapeutic Challenges require no abstract and have no limit for figures and references. Surgical Techniques and Clinicopathologic Correlations

are treated like a full manuscript and require an abstract. All Correspondence and New Instruments should have a standard title page with full length title, running title, and author information. Key words and summary statement should be on the second page. A formal abstract is not required by the journal for Correspondence and New Instruments. A summary statement of 50 words is necessary for publication and indexing and must be included at the time of submission. All pages must be numbered starting with the title page being page one. Each figure must be submitted separately. All color figures will be published in this section at the authors' expense. Authors who submit figures in color do so with the understanding that the figures will be published in color and at their expense.

Manuscripts submitted to **SURGICAL TECHNIQUES** must demonstrate the salient aspects of the technique via video. The video should be submitted in .mp4 format. The video should enhance, but not replace, the technique description in the manuscript. Please see the instructions on Supplemental Digital Content for the details concerning video submission.

The journal accepts manuscripts for consideration as **PHOTO ESSAYS**. These essays include the visual presentation of material where the primary emphasis is on the images. These images can include color images, angiograms, optical coherence tomography, histologic sections, x-rays, ultrasounds, and other studies. The images can be an outstanding presentation of classic findings, atypical findings or new findings, but, the primary emphasis should be on the images. These are not case reports, but rather visual presentation of material as a teaching tool. The images need to be of the highest quality. The accompanying manuscript should be limited to a total of 300 words. A maximum of 5 separate images and 5 references can be included. All figures submitted in color are published in color, at the expense of the author. Please refer to the rest of the author instructions for other requirements for manuscripts submitted to Retina. The Journal doesn't accept Case Reports. Case Reports should be submitted to Retinal Cases & Brief Reports.

Each manuscript will be edited to conform with Editorial Board policy regarding spelling, punctuation, and typographic construction in accordance with "Uniform Requirements for Manuscripts Submitted to Biomedical Journals." As a guide, **The Manual of Style**, prepared by the Scientific Publications Division of the American Medical Association, may be used. When the manuscript has been set in type, proofs will be submitted to the author for approval. These must be returned with the original manuscript and illustrations as indicated on the proofs.

AUTHOR SERVICES

For further guidance AJPM&R and Wolters Kluwer, in partnership with Editage, now offer a unique range of editorial services to help you prepare a submission-ready manuscript. Please visit <http://wkauthorservices.editage.com> to learn more. Services outlined below, as well as others, are available for a nominal cost:

- Premium Editing: Intensive language and structural editing of academic papers to increase chances of journal acceptance.
- Advanced Editing: A complete language, grammar, and terminology check to give you a publication-ready manuscript.

- Translation with Editing: Write your paper in your native language and Wolters Kluwer Author Services will translate it into English, as well as edit it to ensure that it meets international publication standards.
- Plagiarism Check: Helps ensure that your manuscript contains no instances of unintentional plagiarism.
- Artwork Preparation: Save precious time and effort by ensuring that your artwork is viewed favorably by the journal without you having to incur the additional cost of purchasing special graphics software.

After Acceptance

Page proofs and corrections. Corresponding authors will receive electronic page proofs to check the copyedited and typeset article before publication. Portable document format (PDF) files of the typeset pages and support documents (e.g., reprint order form) will be sent to the corresponding author by e-mail. Complete instructions will be provided with the e-mail for downloading and printing the files and for faxing the corrected page proofs to the publisher. Those authors without an e-mail address will receive traditional page proofs. It is the author's responsibility to ensure that there are no errors in the proofs. **Authors must review the proofs thoroughly as errata will not be permitted except for special circumstances.** Changes that have been made to conform to journal style will stand if they do not alter the authors' meaning. Only the most critical changes to the accuracy of the content will be made. Changes that are stylistic or are a reworking of previously accepted material will be disallowed. The publisher reserves the right to deny any changes that do not affect the accuracy of the content. Authors may be charged for alterations to the proofs beyond those required to correct errors or to answer queries. Proofs must be checked carefully and corrections faxed within 24 to 48 hours of receipt, as requested in the cover letter accompanying the page proofs.

Publisher's contact: Email corrected page proofs, reprint order form, and any other related materials to Production Editor Erin Langenfeld at Erin.Langenfeld@wolterskluwer.com.